

PLOCK,ROBERT

Rx #: 7196415

If you have any questions, please feel free to contact your pharmacist at (214)319-0857 or GREER, SHARON S at (800)893-9698

Date: 01/26/2013

Directions : TAKE ONE TABLET BY MOUTH 4 TIMES DAILY AS NEEDED FOR PAIN

IBUPROFEN 600MG TAB

1004

CAUTIONS: THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.