

PLOCK,ROBERT

Rx #: 4486349

If you have any questions, please feel free to contact your pharmacist at (214)319-0857
or GREER, SHARON S at (800)893-9698

Date: 01/26/2013

Directions : TAKE ONE TABLET BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED FOR PAIN NOT
RELIEVED BY MOTRIN

HYDROCO/ACETAMINOP 5-325MG TAB

1004

CAUTIONS: DO NOT TAKE THIS MEDICINE if you are allergic to any ingredient in this medicine. BEFORE USING THIS MEDICINE, check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). DO NOT EXCEED THE RECOMMENDED DOSE, take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS. Make sure that all of your doctors and laboratory personnel know you are taking this medicine. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, emergency care, or surgery, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE DIZZINESS, lightheadedness, or fainting. Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. THIS MEDICINE CONTAINS ACETAMINOPHEN. Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. THIS MEDICINE MAY HARM YOUR LIVER. Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. CONTACT YOUR DOCTOR RIGHT AWAY if you take more than 4,000 mg of acetaminophen per day, even if you feel well. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.