


YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
4702 N Jim Miller Rd
Dallas, TX 75227-
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	ROBERT A PLOCK	DOCTOR	J. KRECK, PBR	DRUG DESCRIPTION
BIRTH DATE	07/26/68			
MEDICATION	HYDROCODONE /ACETAMINOPHEN 10-325 T	PATIENT		
QUANTITY	180	ALLERGIES		
DIRECTIONS	TAKE 1 TO 2 BY MOUTH AS DIRECTED FOR PAIN. NOT TO EXCEED 6 TABLETS PER DAY			



WHITE

FRONT: WATSON 853

INGREDIENT NAME: HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

COMMON USES: This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: THIS MEDICINE CONTAINS ACETAMINOPHEN. Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking sodium oxybate (GHB). ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking anticoagulants (eg, warfarin), buprenorphine, isoniazid, linezolid, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), naltrexone, other narcotic pain medicines (eg, morphine), perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). Ask your doctor if you are unsure if any of your medicines might harm the liver. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), drug or alcohol abuse or dependence, heart problems (eg, cor pulmonale, slow or irregular heartbeat), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you have recently had surgery (eg, stomach or bowel surgery) or are scheduled to have surgery. Tell your doctor if you are in very poor health; are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY or for longer than a few weeks, do not suddenly stop taking it without checking with your doctor. Your doctor may need to gradually lower your dose. SOME OF THESE PRODUCTS CONTAIN SULFITES. Sulfites may cause an allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. STORE THIS MEDICINE at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. Do not store in the bathroom. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you are allergic to any ingredient in this medicine. BEFORE USING THIS MEDICINE, check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). DO NOT EXCEED THE RECOMMENDED DOSE, take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. THIS MEDICINE MAY CAUSE CERTAIN LAB TEST RESULTS. Make sure that all of your doctors and laboratory personnel know you are taking this medicine. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, emergency care, or surgery, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE DIZZINESS, lightheadedness, or fainting. Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. THIS MEDICINE CONTAINS ACETAMINOPHEN. Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. THIS MEDICINE MAY HARM YOUR LIVER. Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. CONTACT YOUR DOCTOR RIGHT AWAY if you take more than 4,000 mg of acetaminophen per day, even if you feel well. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. THIS MEDICINE MAY CAUSE CONSTIPATION. Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. A VERY BAD SKIN REACTION (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

ADDITIONAL INFORMATION: If your symptoms do not improve or if they become worse, contact your doctor. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227
(214)799-7775

RX # 0549165-09015

DATE: 10/16/13

HYDROCODONE /ACETAMINOPHEN 10-325 T

QTY: 180 NO REFILLS - DR. AUTH REQUIRED

New NDC: 00591-2612-05

Retail Price: \$134.89 Your Insurance Saved You: \$124.89

\$ 10.00

J. KRECK, PBR

MFG: WATSON

NJT/NJT/NJT / NJT

PLAN: WPB

GROUP# 01960917

CLAIM REF# 132895830048019998

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227
(214)799-7775

RX # 0549165-09015

DATE: 10/16/13

HYDROCODONE /ACETAMINOPHEN 10-325 T

QTY: 180 NO REFILLS - DR. AUTH REQUIRED

New NDC: 00591-2612-05

Retail Price: \$134.89 Your Insurance Saved You: \$124.89

\$ 10.00

J. KRECK, PBR

MFG: WATSON

NJT/NJT/NJT / NJT

PLAN: WPB

GROUP# 01960917

CLAIM REF# 132895830048019998

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227
PH: (214)388-4951

Customer
Receipt

Pharmacy use only

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227
PH: (214)388-4951

Duplicate
Receipt

WED 5:39PM
New

HYDROCODONE /ACETAMINOPHEN 10-325 T
00591-2612-05
FAST RACK

QTY 180
40 DRAM



WHITE
FRONT: WATSON 853

NJT/NJT/NJT / NJT

Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4702 N Jim Miller Rd
Dallas, TX 75227-
(214)388-4951


IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT ROBERT A PLOCK
BIRTH DATE 07/26/68
MEDICATION CYCLOBENZAPRINE 10MG TABLETS
QUANTITY 90
DIRECTIONS TAKE ONE TABLET BY MOUTH THREE TIMES DAILY

DOCTOR J. KRECK, PBR

DRUG DESCRIPTION

PATIENT ALLERGIES



YELLOW
FRONT: TL177

INGREDIENT NAME: CYCLOBENZAPRINE
(sy-e-kloe-BEN-za-preen)

COMMON USES: This medicine is a muscle relaxant used to treat muscle spasms caused by painful muscle conditions. It should be used along with rest and physical therapy. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine, rasagiline) within the last 14 days. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking bupropion, tramadol, selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine), serotonin norepinephrine reuptake inhibitors (SNRIs) (eg, venlafaxine), anticholinergics (eg, methscopolamine, benztrapine), barbiturates (eg, phenobarbital), meperidine, mibefradil, phenothiazines (eg, chlorpromazine), guanethidine, guanfacine, tricyclic antidepressants (eg, amitriptyline), or verapamil. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor if you have a history of liver problems, heart problems, cerebral palsy, brain or spinal cord disease, stroke, glaucoma, increased pressure in the eye, or trouble urinating. Inform your doctor of any other medical conditions including allergies, pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have an overactive thyroid, moderate to severe liver problems, certain heart problems (eg, irregular heartbeat, congestive heart failure, heart block, conduction problems), or if you have recently had a heart attack. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or are allergic to any ingredient in this product. DO NOT STOP USING THIS MEDICINE without first checking with your doctor. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 2 to 3 weeks without checking with your doctor. If your symptoms (such as pain, tenderness, decreased range of motion) do not improve within 2 to 3 weeks or if they become worse, check with your doctor. THIS MEDICINE MAY CAUSE drowsiness, dizziness, or blurred vision. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Talk with your doctor before you drink alcohol or use medicines that may cause drowsiness (such as sleep aids, muscle

relaxers) while you are using this medicine. This medicine may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. IF YOU EXPERIENCE DRY MOUTH, use sugarless candy or gum, or melt bits of ice in your mouth. If dry mouth continues for more than 2 weeks, contact your dentist or doctor DO NOT BECOME OVERHEATED IN HOT WEATHER or while you are being active; heatstroke may occur. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine especially confusion, hallucinations, and fast or irregular heartbeat. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, drowsiness, dry mouth, fatigue, nausea, nervousness, and stomach pain or upset. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain, confusion, fainting, fast or irregular heartbeat, mental or mood changes, numbness of an arm or a leg, one-sided weakness, seizures, sudden severe stomach pain, severe dizziness or vomiting, speech or vision problems, trouble urinating, or yellowing of the skin or eyes. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; trouble breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include agitation; chest pain; coma; confusion; fast or irregular heartbeat; loss of coordination; hallucinations; seizures; severe drowsiness, dizziness, nausea, or headache; slurred speech; tremor; or vomiting.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227
(214)799-7775

RX # 0549166-09015

DATE: 10/16/13

CYCLOBENZAPRINE 10MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0177-10

Retail Price: \$36.99 Your Insurance Saved You: \$26.99

\$ 10.00

J. KRECK, PBR
MFG: CADISTA
NJT/NJT/NJT/NJT/NJT

PLAN: WPB
GROUP# 01960917
CLAIM REF# 132895831248032999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227-
PH: (214)388-4951

Customer
Receipt

Pharmacy use only

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227
(214)799-7775

RX # 0549166-09015

DATE: 10/16/13

CYCLOBENZAPRINE 10MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0177-10

Retail Price: \$36.99 Your Insurance Saved You: \$26.99

\$ 10.00

J. KRECK, PBR
MFG: CADISTA
NJT/NJT/NJT/NJT/NJT

PLAN: WPB
GROUP# 01960917
CLAIM REF# 132895831248032999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227-
PH: (214)388-4951

Duplicate
Receipt

WED 5:39PM
New

CYCLOBENZAPRINE 10MG TABLETS
59746-0177-10
FAST RACK

QTY 90
10 DRAM



YELLOW
FRONT: TL177

NJT/NJT/NJT/NJT/NJT

Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

Walgreens

#09015 4702 N JIM MILLER RD
DALLAS, TX 75227
214-388-4951

805 6981 0091 10/16/2013 8:05 PM

FSA RX 0549165	10.00
CLM RF# 132895830048019998	
FSA RX 0549166	10.00
CLM RF# 132895831248032999	

REWARDS BONUS EARNED 1000

TOTAL	20.00
VISA ACCT 2677	20.00
CHANGE	.00

TOTAL FSA ITEMS	0.00
TOTAL RX ITEMS	20.00
TOTAL FSA AND RX ITEMS	20.00

APPROVED FSA/HRA AMOUNT 0.00

THANK YOU FOR SHOPPING AT WALGREENS

REDEEM AT THE REGISTER WITH AS FEW AS
5000 POINTS. RESTRICTIONS APPLY, SEE
PROGRAM RULES FOR DETAILS. PLEASE GO
TO WALGREENS.COM/BALANCE.

RFN# 0901-5916-9812-1310-1603



Q