

# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location


4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT ROBERT A PLOCK  
BIRTH DATE 07/26/68  
MEDICATION METHYLPREDNISOLONE 4MG DOSPAK 21'S  
QUANTITY 21  
DIRECTIONS TAKE AS DIRECTED

DOCTOR L. JACKSON, PA  
PATIENT ALLERGIES

### DRUG DESCRIPTION



WHITE  
FRONT: TL 001

**INGREDIENT NAME:** METHYLPREDNISOLONE  
(meth-ill-pred-NISS-oh-lone)

**COMMON USES:** This medicine is a corticosteroid used to treat severe allergies, arthritis, asthma, and skin conditions. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking mifepristone. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking aspirin, barbiturates, cyclosporine, licorice (glycyrrhiza), macrolide antibiotics (e.g., erythromycin), indomethacin, estrogen (e.g., birth control pills), ketoconazole, rifampin, troleandomycin, or medicine for seizures. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a fungal infection in the body. Inform your doctor of any medical conditions including heart disease, liver disease, kidney disease, HIV infection, high blood pressure, peptic ulcer, osteoporosis, myasthenia gravis, herpes eye infection, low levels of thyroid hormone (hypothyroidism), diabetes, tuberculosis, mental/mood disorders, seizures, recent infection, allergies, pregnancy, or breast-feeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. The directions for taking this medicine properly are also on the package. This medicine may be taken on an empty stomach or with food. Taking it with food may help to decrease the chance of stomach upset. Ask your doctor or pharmacist any questions you may have about how you should take this medicine. STORE THIS MEDICINE at room temperature, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking 1 dose daily, take it as soon as possible. If you do not remember until the next day, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once. For all other dosing regimens: if you miss a dose, contact your doctor or pharmacist promptly. Your dosing schedule may need to be adjusted.

**CAUTIONS:** DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. Stopping this medicine suddenly may cause serious side effects. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are using this medicine. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. CORTICOSTEROIDS MAY LOWER YOUR ABILITY TO FIGHT INFECTION or may cause infections to be more severe. The risk may be greater if you take them for a long time. Avoid contact with people who have infections such as colds, chickenpox, measles, or tuberculosis (TB). Tell your doctor right away if you have contact with anyone who has these infections. Tell your doctor if you develop symptoms such as fever, sore throat, rash, or chills. Check with your doctor if you have questions about your risk of infection. CHECK WITH YOUR DOCTOR BEFORE HAVING IMMUNIZATIONS (VACCINATIONS) while you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either

prescription or over-the-counter, check with your doctor or pharmacist. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. IF YOU HAVE DIABETES, this medicine may affect your blood sugar. Check blood sugar levels closely and ask your doctor before adjusting the dose of your diabetes medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while you are taking this medicine include dizziness; facial flushing; feeling of whirling motion; headache; or increased sweating. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience bloody, black, or tarry stools; changes in body fat; changes in menstrual periods; chest pain; fainting; fever, chills, or sore throat; increased hunger, thirst, or urination; mental or mood changes; muscle pain, weakness, or wasting; seizures; severe nausea or vomiting; slow wound healing; stomach pain; sudden severe dizziness or headache; swelling of feet or legs; tendon, bone, or joint pain; thinning of the skin; unusual bruising or bleeding; unusual skin sensations; unusual weight gain; vision changes or other eye problems; or vomit that looks like coffee grounds. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; or swelling of the mouth, face, lips, or tongue. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately.

**ADDITIONAL INFORMATION:** Patients taking this medicine for long-term therapy should carry a warning card or medical ID bracelet that identifies the possible need of supplemental steroids during periods of stress. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0643548-09015

DATE: 01/14/15

#### METHYLPREDNISOLONE 4MG DOSPAK 21'S

QTY: 21 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0001-03

Retail Price: \$32.99 Your Insurance Saved You: \$12.99

\$ 20.00

L. JACKSON, PA  
MFG: CADISTA  
RRL/PDR/PDR/ /PDR

PLAN: TXBC  
GROUP# 001740  
CLAIM REF# 150143014073017999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951

Customer  
Receipt

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0643548-09015

DATE: 01/14/15

#### METHYLPREDNISOLONE 4MG DOSPAK 21'S

QTY: 21 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0001-03

Retail Price: \$32.99 Your Insurance Saved You: \$12.99

\$ 20.00

L. JACKSON, PA  
MFG: CADISTA  
RRL/PDR/PDR/ /PDR

PLAN: TXBC  
GROUP# 001740  
CLAIM REF# 150143014073017999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

WED 9:51AM  
New

METHYLPREDNISOLONE 4MG DOSPAK 21'S  
59746-0001-03  
FAST RACK

QTY 21



WHITE  
FRONT: TL 001

RRL/PDR/PDR/ /PDR

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382